

Section 3.1

Math 231

Hope College

Vector Spaces

A **vector space over** \mathbb{R} is a set V of objects (called **vectors**), together with two operations, addition and scalar multiplication, which satisfy the following:

- 1 V is closed under addition.
- 2 V is closed under scalar multiplication.
- 3 For all $\mathbf{x}, \mathbf{y} \in V$, we have $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$.
- 4 For all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$, we have $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$.
- 5 There exists $\mathbf{0} \in V$ such that for all $\mathbf{x} \in V$, we have $\mathbf{x} + \mathbf{0} = \mathbf{x}$. (The vector $\mathbf{0}$ is called a **zero vector** for V .)
- 6 For each $\mathbf{x} \in V$, there exists $\mathbf{y} \in V$ such that $\mathbf{x} + \mathbf{y} = \mathbf{0}$. (\mathbf{y} is called an **additive inverse** of \mathbf{x} .)
- 7 For all $\mathbf{x} \in V$, we have $1\mathbf{x} = \mathbf{x}$.
- 8 For all $\alpha, \beta \in \mathbb{R}$ and all $\mathbf{x} \in V$, we have $(\alpha\beta)\mathbf{x} = \alpha(\beta\mathbf{x})$.
- 9 For all $\alpha \in \mathbb{R}$ and all $\mathbf{x}, \mathbf{y} \in V$, we have $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}$.
- 10 For all $\alpha, \beta \in \mathbb{R}$ and all $\mathbf{x} \in V$, we have $(\alpha + \beta)\mathbf{x} = \alpha\mathbf{x} + \beta\mathbf{x}$.



Examples of Vector Spaces

- 1 For all n , \mathbb{R}^n is a vector space.
- 2 For all m, n , $M_{m,n}(\mathbb{R})$ is a vector space.
- 3 The set $\mathcal{P}(\mathbb{R})$ of all polynomials in one variable x with real coefficients is a vector space.
- 4 The set $\mathcal{P}_n(\mathbb{R})$ of all polynomials of degree at most n in one variable x with real coefficients is a vector space.
- 5 The set $\mathcal{F}(\mathbb{R})$ of all functions from \mathbb{R} to \mathbb{R} is a vector space.

Examples of Vector Spaces

- 1 For all n , \mathbb{R}^n is a vector space.
- 2 For all m, n , $M_{m,n}(\mathbb{R})$ is a vector space.
- 3 The set $\mathcal{P}(\mathbb{R})$ of all polynomials in one variable x with real coefficients is a vector space.
- 4 The set $\mathcal{P}_n(\mathbb{R})$ of all polynomials of degree at most n in one variable x with real coefficients is a vector space.
- 5 The set $\mathcal{F}(\mathbb{R})$ of all functions from \mathbb{R} to \mathbb{R} is a vector space.

Examples of Vector Spaces

- 1 For all n , \mathbb{R}^n is a vector space.
- 2 For all m, n , $M_{m,n}(\mathbb{R})$ is a vector space.
- 3 The set $\mathcal{P}(\mathbb{R})$ of all polynomials in one variable x with real coefficients is a vector space.
- 4 The set $\mathcal{P}_n(\mathbb{R})$ of all polynomials of degree at most n in one variable x with real coefficients is a vector space.
- 5 The set $\mathcal{F}(\mathbb{R})$ of all functions from \mathbb{R} to \mathbb{R} is a vector space.

Examples of Vector Spaces

- 1 For all n , \mathbb{R}^n is a vector space.
- 2 For all m, n , $M_{m,n}(\mathbb{R})$ is a vector space.
- 3 The set $\mathcal{P}(\mathbb{R})$ of all polynomials in one variable x with real coefficients is a vector space.
- 4 The set $\mathcal{P}_n(\mathbb{R})$ of all polynomials of degree at most n in one variable x with real coefficients is a vector space.
- 5 The set $\mathcal{F}(\mathbb{R})$ of all functions from \mathbb{R} to \mathbb{R} is a vector space.

Examples of Vector Spaces

- 1 For all n , \mathbb{R}^n is a vector space.
- 2 For all m, n , $M_{m,n}(\mathbb{R})$ is a vector space.
- 3 The set $\mathcal{P}(\mathbb{R})$ of all polynomials in one variable x with real coefficients is a vector space.
- 4 The set $\mathcal{P}_n(\mathbb{R})$ of all polynomials of degree at most n in one variable x with real coefficients is a vector space.
- 5 The set $\mathcal{F}(\mathbb{R})$ of all functions from \mathbb{R} to \mathbb{R} is a vector space.

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .

Properties of Vector Spaces

Theorem 3.7: Let V be a vector space.

- 1 The zero vector $\mathbf{0}$ is unique.
- 2 Given $\mathbf{x} \in V$, its additive inverse is unique.
- 3 Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$. If $\mathbf{x} + \mathbf{z} = \mathbf{y} + \mathbf{z}$, then $\mathbf{x} = \mathbf{y}$.
- 4 For all $\mathbf{x} \in V$, we have $0\mathbf{x} = \mathbf{0}$.
- 5 For all $\alpha \in \mathbb{R}$, we have $\alpha\mathbf{0} = \mathbf{0}$.
- 6 For all $\mathbf{x} \in V$, the vector $(-1)\mathbf{x}$ is the additive inverse of \mathbf{x} .