

Section 4.1

Math 231

Hope College

Function, Domain, and Image

Let X and Y be sets.

- 1 A **function** f from X to Y is a rule which assigns one and only one element $f(x) \in Y$ to each element $x \in X$.
- 2 We denote a function f from X to Y by $f : X \rightarrow Y$.
- 3 The set X is called the **domain** of f . We write $X = \text{dom}(f)$.
- 4 The subset $\{y \in Y \mid y = f(x) \text{ for some } x \in X\}$ of Y is called the **range** or the **image** of f . We write the image of f as $\text{im}(f)$.

Surjective, Injective, and Bijective

Let X and Y be sets, and let $f : X \rightarrow Y$ be a function.

- 1 f is called **surjective** if $\text{im}(f) = Y$.
- 2 f is called **injective** if $x_1, x_2 \in X$ and $x_1 \neq x_2$ implies that $f(x_1) \neq f(x_2)$. Thus, f is injective if distinct inputs have distinct outputs.
- 3 The statement that f is injective is equivalent to

for all $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies that $x_1 = x_2$.
- 4 f is called **bijective** if it is both injective and surjective.

Identity, Composition, Inverse, and Invertible

Let X , Y and Z be sets.

- 1 The **identity function** $\text{id}_X : X \rightarrow X$ is the function defined by $\text{id}_X(x) = x$ for all $x \in X$.
- 2 Given functions $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, the **composition** $g \circ f$ is the function $g \circ f : X \rightarrow Z$ defined by $g \circ f(x) = g(f(x))$ for all $x \in X$.
- 3 Given a function $f : X \rightarrow Y$, an **inverse function** for f is a function $f^{-1} : Y \rightarrow X$ such that $f^{-1} \circ f = \text{id}_X$ and $f \circ f^{-1} = \text{id}_Y$.
- 4 If a function f has an inverse function f^{-1} , we say that f is an **invertible** function.

Theorem 4.3: A function is invertible if and only if it is bijective.

Linear Transformations

- Let $f : V \rightarrow W$ be a function from a vector space V to a vector space W . Then f is called a **linear transformation** if the following two conditions are satisfied:
 - 1 For all $\mathbf{x}, \mathbf{y} \in V$, we have $f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$.
 - 2 For all $\mathbf{x} \in V$ and $\alpha \in \mathbb{R}$, we have $f(\alpha\mathbf{x}) = \alpha f(\mathbf{x})$.

- **Theorem 4.7:** Let A be an $m \times n$ matrix. Then A defines a linear transformation $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $f(\vec{\mathbf{x}}) = A\vec{\mathbf{x}}$.

Linear Transformations

- Let $f : V \rightarrow W$ be a function from a vector space V to a vector space W . Then f is called a **linear transformation** if the following two conditions are satisfied:
 - 1 For all $\mathbf{x}, \mathbf{y} \in V$, we have $f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$.
 - 2 For all $\mathbf{x} \in V$ and $\alpha \in \mathbb{R}$, we have $f(\alpha\mathbf{x}) = \alpha f(\mathbf{x})$.

- **Theorem 4.7:** Let A be an $m \times n$ matrix. Then A defines a linear transformation $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $f(\vec{\mathbf{x}}) = A\vec{\mathbf{x}}$.

Linear Transformations

- Let $f : V \rightarrow W$ be a function from a vector space V to a vector space W . Then f is called a **linear transformation** if the following two conditions are satisfied:
 - For all $\mathbf{x}, \mathbf{y} \in V$, we have $f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$.
 - For all $\mathbf{x} \in V$ and $\alpha \in \mathbb{R}$, we have $f(\alpha\mathbf{x}) = \alpha f(\mathbf{x})$.
- Theorem 4.7:** Let A be an $m \times n$ matrix. Then A defines a linear transformation $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $f(\vec{\mathbf{x}}) = A\vec{\mathbf{x}}$.

Moreover, given any linear transformation $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there is an $m \times n$ matrix A such that $f(\vec{\mathbf{x}}) = A\vec{\mathbf{x}}$ for all $\vec{\mathbf{x}} \in \mathbb{R}^n$.